


The listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1. (Previously Presented) A method of treatment of a disorder selected from the group consisting of obesity, excessive food intake, and smoking craving comprising administration to a subject in need of such treatment an effective dose of a compound of formula (I):



(I)

wherein:

R<sup>1</sup> is aryl;

R<sup>2</sup> is H, alkyl or aryl; and

R<sup>3</sup> is hydrogen or alkyl;

or a pharmaceutically acceptable salt thereof.

2. (Previously Presented) A method according to claim 1 wherein R<sup>1</sup> is a substituted or unsubstituted phenyl or naphthyl.

3. (Previously Presented) A method according to claim 1 wherein R<sup>1</sup> has 1, 2 or 3 substituent groups.

4. (Previously Presented) A method according to claim 1 wherein R<sup>1</sup> is chlorophenyl or (trifluoromethyl)phenyl.

5. (Previously Presented) A method according to claim 1 wherein R<sup>2</sup> is aryl.

6. (Previously Presented) A method according to claim 5 wherein R<sup>3</sup> is alkyl.

7. (Previously Presented) A method according to claim 1 wherein the compound is selected from:  
3-(bis(4-chlorophenyl)methoxy)-N-(2-propenyl)azetidine-1-carboxamide; and  
(R)-3-(bis(4-chlorophenyl)methoxy)-N-(2-hydroxypropyl)azetidine-1-carboxamide.
8. (Previously Presented) A method according to claim 1 wherein the dose further comprises a pharmaceutically acceptable carrier.
9. (Canceled)
10. (Canceled)
11. (Previously Presented) A method according to claim 1 wherein the disorder is obesity.
12. (Currently Amended) A method according to claim 1 ~~for~~ wherein the disorder is smoking craving.
13. (Previously Presented) A method according to claim 1 wherein the disorder is excessive food intake.
14. (Previously Presented) A method according to claim 2 wherein R<sup>1</sup> is a substituted or unsubstituted phenyl or naphthyl.
15. (Previously Presented) A method according to claim 2 wherein R<sup>1</sup> has 1, 2 or 3 substituent groups.
16. (Previously Presented) A method according to claim 2 wherein R<sup>1</sup> is chlorophenyl or (trifluoromethyl)phenyl.
17. (Previously Presented) A method according to claim 1 wherein R<sup>2</sup> is aryl.
18. (Canceled)

19. (Previously Presented) A method according to claim 7 wherein the disorder is obesity.

20. (Currently Amended) A method according to claim 7 ~~for~~ wherein the disorder is smoking craving.

21. (Previously Presented) A method according to claim 7 wherein the disorder is excessive food intake.